



**RECALL: Elimination reactions**

$-\text{OH}$  and  $-\text{OCH}_3$  are bad leaving groups and so these reactions would not occur spontaneously without an acid catalyst.


To determine whether an elimination can occur, ask yourself three questions:




1. Is there a good leaving group present?  
Eg. Yes, Br is a good leaving group
2. Is there a hydrogen on the carbon next to the carbon containing the leaving group?  
Eg. Yes, on the bridge-heads on either side of the carbon containing the Br.
3. Is Bredt's Rule being followed?  
Eg. No, if a double bond was being formed, it would be at a bridge-head and Bredt's rule states that a double bond cannot be formed at a bridge-head if the rings are small and all bridges  $>0$ . (double bond too strained)

Substitution would likely not occur either as the electrophilic site is hindered (tertiary carbon).



## Addition Reactions




For alternate regiochemistry (addition of Br onto the less substituted carbon) need dialkyl peroxide






## Examples of peroxides



## Radical mechanism



Example without HBr – a polymerization reaction occurs



shorthand notation for polypropylene

Note more stable radical (in this case tertiary) is always formed by addition onto double bond

There is a termination step (not shown) that ends this polymerization. It requires combination of 2 radicals. It could be two growing chain radicals meeting or it could be from peroxide. If less peroxide is used the polymer chain will be longer.

## Polymers

Poly = many


Meros = parts

## Examples of Biopolymers

1. Polysaccharides  
- polymers of sugars
2. Proteins and peptides  
- polymers of amino acids
3. Nucleic acid polymers (DNA and RNA)  
- polymers of nucleotides
4. Fats and polyketides  
- polymers of fatty acids
5. Polyisoprenoids/ terpenoids  
- polymers of isoprene


## Polymer formation

### Teflon




Many polymers degrade into their components if heated enough, and can further decompose.


### Polyethylene



## Polyacrylonitrile




Found in carpets




Polyacrylonitrile can form HCN if it is heated to decomposition.

## Polyvinyl chloride




Such polymers containing chloride can form HCl if decomposed.

## Copolymers




Copolymers are composed of two different subunits.

### Lucite




### Polystyrene




phenyl group is Ph

### Example: Mechanism of polystyrene formation



### Short-hand for mechanism of polystyrene formation



Divinyl benzene can be added as a cross-linker so chains link on both of its double bonds  
This make the copolymer more solid (as you encounter in many products) – typically  
about one part in 100 to one part in 6 of divinylbenzene may be added

