

REACTIONS OF CARBOXYLIC ACIDS & DERIVATIVES

Important Common Names of Carboxylic Acids Include:

Systematic Name	Common Name	Formula
methanoic acid	formic acid	HCO_2H
ethanoic acid	acetic acid	$\text{CH}_3\text{CO}_2\text{H}$
propanoic acid	propionic acid	$\text{CH}_3\text{CH}_2\text{CO}_2\text{H}$
butanoic acid	butyric acid	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$
pentanoic acid	valeric acid	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$
benzoic acid	benzoic acid	PhCO_2H
ethanedioic acid	oxalic acid	$\text{HO}_2\text{C}-\text{CO}_2\text{H}$
propanedioic acid	malonic acid	$\text{HO}_2\text{CCH}_2\text{CO}_2\text{H}$
butanedioic acid	succinic acid	$\text{HO}_2\text{CCH}_2\text{CH}_2\text{CO}_2\text{H}$
pentanedioic acid	glutaric acid	$\text{HO}_2\text{CCH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$
hexanedioic acid	adipic acid	$\text{HO}_2\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{H}$

Reactions of carboxylic acids and their derivatives (acyl halides, anhydrides, esters, amides) resemble those of ketones and aldehydes, but replacement (substitution) of an electronegative group on the carbonyl is the common extra feature. Most reactions follow a pattern:


1. Addition to carbonyl by strong (irreversible) or weak (reversible) nucleophiles with expulsion of an electronegative group.

⇒ The strong nucleophiles (hydride and alkyl anion) usually add a second time to the aldehyde or ketone that has been generated by the first addition


2. Reaction at alpha carbon with an electrophile via enolate

1. Addition to Carbonyl

- In the first step, nucleophiles generally attack the carbonyl of carboxylic acid derivatives (e.g. acid chlorides, anhydrides, esters, amides) in the same way as with ketones and aldehydes.
- However, the presence of an electron withdrawing atom directly attached to the former carbonyl carbon allows its expulsion to generate a new carbonyl derivative in which the nucleophile has replaced the heteroatom group (Y).
- In the case of interconversions of carboxylic acid derivatives with heteroatom (weak) nucleophiles, this gives the final product in the box.




- If the nucleophile is very strong, such as R^- (alkyl anion from RLi or RMgX) or H^- (hydride anion from LiAlH_4 or NaBH_4), there is no H^+ present (the positive ion is a metal) and the product in the box is a ketone or aldehyde which will react with a second mole of nucleophile (see ketone reactions) to ultimately generate an alcohol (in the circle).
- Amides are an exception and give amines with LiAlH_4 or NaBH_4 .
- In order to decide whether a reaction with a heteroatom nucleophile will work, the scheme shown below is useful:




- In general, carboxylic acid derivatives that are more reactive can be spontaneously converted into less reactive ones by treatment with the appropriate nucleophilic species.
- The exceptions are carboxylic acids which will generally react with alcohols to form esters only under acidic conditions because under basic conditions the formation of unreactive salts is extremely rapid.
- The general reaction for formation of carboxylic acid derivatives is:

Under Basic Conditions:



Under Acidic Conditions:



2. Reaction at alpha Carbon

- The general features of this type of reaction are analogous to the corresponding reactions of aldehydes and ketones. Of course the products will now be carboxylic acid derivatives.
- This type of reaction is particularly useful with esters.
- The base must be chosen carefully such that it does not act as a nucleophile and attack the carbonyl in a "Type 1" reaction. For esters, LDA (lithium diisopropylamide) at low temperature is especially useful as a base.

